

## SUPPLEMENTARY TABLES

**Supplementary Table 1. The clinicopathologic characteristics of enrolled patients for circRNA CHIP.**

	<b>Age</b>	<b>Gender</b>	<b>Pathological types</b>	<b>TNM stage</b>	<b>Pathological stage</b>
Patient 1	67	Female	Squamous cell carcinoma	IV	Low
Patient 2	54	Male	Squamous cell carcinoma	III	middle
Patient 3	60	Male	Squamous cell carcinoma	IV	Low
Patient 4	68	Female	Squamous cell carcinoma	IV	middle
Patient 5	59	Female	Squamous cell carcinoma	IV	Low

**Supplementary Table 2. Patients' clinicopathologic characteristics (N=58).**

<b>Characteristics</b>	<b>Cases (n)</b>
<i>Age (Median /years)</i>	65(42-75) y
<i>Gender</i>	
Male	38(65.52)
Female	20(34.48)
<i>Pathological types</i>	
Adenocarcinoma	0(0.00)
Squamous cell carcinoma	58(100.00)
<i>TNM stage</i>	
II/ III	25(43.11)
IV	33(56.89)
<i>Pathological stage</i>	
High and middle	22(37.93)
Low	36(52.07)
<i>EGFR mutation site</i>	
19del/L858R	51(87.93)
others	7(12.07)